

**Page 18, line 16, paragraph 5:** Please amend as follows:

B1  
Figs. 5A-5U: Gene sequences of the synthetic V heavy chain consensus genes. The corresponding amino acid sequences (see Figs. 2E-2G) as well as the unique cleavage sites are also shown.

**Page 18, line 19, paragraph 6:** Please amend as follows:

Figs. 6A-6G: Oligonucleotides used for construction of the consensus genes. The oligos are named according to the corresponding consensus gene, e.g. the gene V $\kappa$ 1 was constructed using the six oligonucleotides O1K1 to O1K6. The oligonucleotides used for synthesizing the genes encoding the constant domains C $\kappa$  (OCLK1 to 8) and CH1 (OCH1 to 8) are also shown.

**Page 18, line 25, paragraph 7:** Please amend as follows:

Figs. 7A-7D: Sequences of the synthetic genes encoding the constant domains C $\kappa$  (7A-7B) and CH1 (7C-7D). The corresponding amino acid sequences as well as unique cleavage sites introduced in these genes are also shown.

**Page 18, line 28, paragraph 8:** Please amend as follows:

Figs. 7E-7H: Functional map and sequence of module M24 comprising the synthetic C $\lambda$  gene segment (huCL lambda).

**Page 18, line 30, paragraph 9:** Please amend as follows:

Figs. 7I-7J: Oligonucleotides used for synthesis of module M24.

**Page 18, line 31, paragraph 10:** Please amend as follows:

Figs. 8A-8E: Sequence and restriction map of the synthetic gene encoding the consensus single-chain fragment VH3-V $\kappa$ 2. The signal sequence (amino acids 1 to 21) was derived from the *E. coli* phoA gene (Skerra & Plückthun, 1988). Between the phoA signal sequence and the VH3 domain, a short sequence stretch encoding 4 amino acid residues (amino acid 22 to 25) has been inserted in order to allow detection of the single-chain fragment in Western blot or ELISA using the monoclonal antibody M1

B1  
(Knappik & Plückthun, 1994). The last 6 basepairs of the sequence were introduced for cloning purposes (EcoRI site).

Page 19, line 14, paragraph 3: Please amend as follows:

B2  
SUB  
C2  
Figs. 10A-10B: Sequencing results of independent clones from the initial library, translated into the corresponding amino acid sequences. (A) Amino acid sequence of the VH3 consensus heavy chain CDR3 (position 93 to 102, Kabat numbering). (B) Amino acid sequences of 12 clones of the 10-mer library. (C) Amino acid sequences of 11 clones of the 15-mer library, \*: single base deletion.

Page 20, line 35, paragraph 10: Please amend as follows:

B3  
Fig. 25A: Schematic representation of the modular pCAL vector system.

Page 20, line 36, paragraph 11: Please amend as follows:

B4  
Figs. 25B-25C: List of restriction sites already used in or suitable for the modular HuCAL genes and pCAL vector system.

Page 20, line 38, paragraph 12: Please amend as follows:

Figs. 26A-26D: List of the modular vector elements for the pCAL vector series: shown are only those restriction sites which are part of the modular system.

Page 21, line 1, paragraph 1: Please amend as follows:

SUB  
C5  
Figs. 27A-27B: Functional map and sequence of the multi-cloning site module (MCS).

Page 21, line 2, paragraph 2: Please amend as follows:

Figs. 28A-28G: Functional map and sequence of the pMCS cloning vector series.

Page 21, line 3, paragraph 3: Please amend as follows:

Figs. 29A-29B: Functional map and sequence of the pCAL module M1 (see Figs. 26A-26D).

B4 [Page 21, line 4, paragraph 4: Please amend as follows:]

Figs. 30A-30C: Functional map and sequence of the pCAL module M7-III (see Figs. 26A-26D).

[Page 21, line 5, paragraph 5: Please amend as follows:]

Figs. 31A-31B: Functional map and sequence of the pCAL module M9-II (see Figs. 26A-26D).

[Page 21, line 6, paragraph 6: Please amend as follows:]

Figs. 32A-32C: Functional map and sequence of the pCAL module M11-II (see Figs. 26A-26D).

[Page 21, line 7, paragraph 7: Please amend as follows:]

Figs. 33A-33D: Functional map and sequence of the pCAL module M14-Ext2 (see Figs. 26A-26D).

[Page 21, line 9, paragraph 8: Please amend as follows:]

Figs. 34A-34D: Functional map and sequence of the pCAL module M17 (see Figs. 26A-26D).

[Page 21, line 10, paragraph 9: Please amend as follows:]

Figs. 35A-35I: Functional map and sequence module vector pCAL4.

B4 [Page 21, line 11, paragraph 10: Please amend as follows:]

Figs. 35J-35XXX: Functional maps and sequences of additional pCAL modules (M2, M3, M7I, M7II, M8, M10II, M11II, M12, M13, M19, M20, M21, M41) and of low-copy number plasmid vectors (pCALO1 to pCALO3).

[Page 21, line 14, paragraph 11: Please amend as follows:]

Figs. 35YYY-35CCCC: List of oligonucleotides and primers used for synthesis of pCAL vector modules.